

Update on School (K – 12) and Childcare Facilities: Interim CDC Guidance in Response to Human Infections with the Novel Influenza A (H1N1) Virus

Taken from a CDC document revised May 4, 2009, 8:30PM EDT

Background

This document provides updated interim guidance for schools and childcare facilities regarding the prevention of the spread of novel influenza A (H1N1) virus.

Initial cases of novel influenza A (H1N1) in the United States included school-aged students and were associated with travel to Mexico and school-based outbreaks. Based on initial information, CDC recommended consideration of school closure as an option to lessen the risk of infection with this novel influenza virus in order to protect students, staff, parents and other caregivers from a potentially severe disease as well as limit spread into the community.

New information on disease severity warrant revision of the school closure guidance. Most U.S. cases have not been severe and are comparable in severity to seasonal influenza. CDC and local and state health officials will continue to closely monitor the severity and spread of this novel H1N1 influenza outbreak.

At this time, CDC recommends the primary means to reduce spread of influenza in schools focus on early identification of ill students and staff, staying home when ill, and good cough and hand hygiene etiquette. Decisions about school closure will be at the discretion of local authorities based on local considerations, including public concern and the impact of school absenteeism and staffing shortages.

Recommendations

- School closure **is not advised** for a suspected or confirmed case of novel influenza A (H1N1) and, in general, is not advised unless there is a magnitude of faculty or student absenteeism that interferes with the school's ability to function.
- Students, faculty or staff with influenza-like illness (fever with a cough or sore throat) should stay home and not attend school or go into the community except to seek medical care for at least 7 days even if symptoms resolve sooner.
- Students, faculty and staff who appear to have an influenza-like illness at arrival or become ill during the school day should be isolated promptly in a room separate from other students and sent home.
- Parents and guardians should monitor their school-aged children, and faculty and staff should self-monitor every morning for symptoms of influenza-like illness.
- Ill students should not attend afterschool programs or congregate in settings other than school.
- School administrators should communicate regularly with local public health officials to obtain guidance about reporting of influenza-like illnesses in the school.
- Students, faculty and staff should stringently follow sanitary measures to reduce the spread of influenza, including covering their nose and mouth with a tissue when coughing or sneezing (or coughing or sneezing into their sleeve if a tissue isn't available), frequently washing hands with soap and water, or using hand sanitizer if hand washing with soap and water is not possible.

Further guidance can be found in:

Questions and Answers About Novel H1N1 Flu at http://www.cdc.gov/h1n1flu/swineflu_you.htm

What to Do If You Get Flu-Like Symptoms at <http://www.cdc.gov/h1n1flu/sick.htm>

Interim Guidance for H1N1 Flu: Taking Care of a Sick Person in Your Home at http://www.cdc.gov/h1n1flu/guidance_homecare.htm