Student Name:	_ Age:	Grade:
School:	Homeroom Teacher:	
Parent/Guardian:	Phone(H)	(W)
Parent/Guardian:	Phone(H)	(W)
Alternate Emergency Contact;	Phone(H)	(W)
Know Asthma Triggers: Please check the boxes to identify what can ca Exercise Respiratory/viral infections Odors/fumes/ Pollens Animals/dander Dust/dust mit Temperature/weather – humidity, cold air, etc. Pesticides Other- please list:	/smoke Mold/ es Grasse Food -	mildew es/trees - Please list below
Know Allergy/Intolerance: Please check those which apply and describe whe contact with the allergen.	nat happens when your	child eats or comes into
Peanuts Tree Nuts Fish/Shellfish ggs Soy Wheat Milk Medication Latex Insect stings Other Notice: If your child has been prescribed epinephrine (e.g. EpiPen) for an alle school. If your student requires a special diet to limit or eliminate foods, your se "Medical Statement tor Students Requiring special Meals".	ergy, it is also necessary	to provide epinephrine at
<u>Daily Medications:</u> Please list daily medications used at home and/or to be administered at school. Medication Name Amount/Dose When administered		
I understand that all medications to be administered at school must be provided by the parent/guardian.		
Parent signature:	D:	ate:
Reviewed by school nurse/nurse designee:	Da	ate:



Student Name: _____ Date Of Birth: _______ Exercise Pre-Treatment: Administer inhaler (2 Inhalations) 15-30 minutes prior to exercise. (e.g. PE, recess, etc). Albuterol HFA inhaler (Proventil, Ventolin, ProAir) Use inhaler with spacer/valved holding chamber Levalbuterol (Xopenex HFA) May carry & self-administer inhaler (MD) Pirbuterol inhaler (Maxair) Other: **Asthma Treatment Anaphylaxis Treatment** Give quick relief medication when student experiences Give epinephrine when student experiences allergy asthma symptoms, such as coughing, wheezing or tight chest Symptoms, such as tongue swelling, throat closing, change in voice, faintness, difficulty breathing (chest Abuterol HFA (Proventil, Ventolin, ProAir) 2 inhalations or neck "sucking in), lips or fingernails turning blue, Levalbuterol (Xopenex HFA) 2 inhalations or trouble talking (shortness of breath). Pirbuterol (Maxair) 2 inhalations EpiPen® 0.3 mg Use inhaler with spacer/valved holding chamber EpiPen® jr. 0.15 mg May cary & self-administer inhaler (MD) TwinjectTM 0.3 mg Albuterol inhaled by nebulizer (Proventil, Ventolin, Twinject TM 0.15 mg AccuNeb) .63 mg/3 mL 1.25 mg/3 mL 2.5 mg/3 mL Adrenaclick® 0.3 mg Levalbuterol inhaled by nebulizer (Xopenex) Adrenaclick® 0.15 mg 0.3 mg/3 mL 0.63 mg/3 mL 1.25 mg/3 mL Other: Other: May carry & self-administer epinephrine Closely Observe the Student after CALL 911 After Giving Epinephrine, Closely Giving Quick Relief Medication **Observe the Student** If, after 10 minutes: Notify parent/guardian immediately Symptoms are improved, student may return to Even if student improves, the student Classroom after notifying parent/guardian Should be observed for recurrent No improvement in symptoms, repeat the treatment Symptoms of anaphylaxis in an emergency and notify parent/guardian immediately medical facility If student continues to worsen CALL 911 and If student does not improve or continues to Initiate the Richmond County Schools' Emergency worsen, consider a second dose of Response to LifeThreatening Asthma or Systemic epinephrine and initiate Life Threatening Allergic Reactions (Anaphylaxis) Protocol Allergic Reaction Protocol In this student has a medical history of asthma and/or anaphylaxis and I have reviewed the use of the above-listed medication(s). If Medications are self-administered; the school staff must be notified. Additional information: (i.e asthma triggers, allergens) Physician name: (please print) hysician Signature: Parent Signature: Reviewed by school nurse/nurse designee: